

PRESSEMITTEILUNG

Current loss tracked down by magnetic fingerprint

HZB researchers solve the case of lost current in organic solar cells

Berlin, 26. 10. 2010

Conventional solar cells made from crystalline silicon are difficult and energy-intensive to manufacture. Organic solar cells are cheaper, but have always produced less electricity. Why this is so has never been fully explained. Now, a method developed by HZB researchers reveals that current flow inside a solar cell can be affected by the spin of the charge-carrying particles.

Further informations:

Dr. Klaus Lips
Institut Silizium-Photovoltaik
Tel.: +49 (0)30-8062-41353
lips@helmholtz-berlin.de

Scientists have been working on organic solar cells for about a decade. Their manufacture is environmentally friendly and they can be applied to all kinds of materials, such as plastic film, for instance. The trouble is, they only yield a fifth of the electrical energy that silicon solar cells do, with most of the electrical current trickling away into the material instead.

Dr. Alexander Schnegg
Institut Silizium-Photovoltaik
Tel.: +49 (0)30-8062-41373
alexander.schnegg@helmholtz-berlin.de

Scientists at Helmholtz-Zentrum Berlin (HZB) have developed a method that uses the magnetic fingerprint of the charge-carrying particles to reveal exactly how electricity is being lost. They did so by cleverly manipulating the magnetic properties of these particles. Together with Scottish researchers, they have published their findings in *Physical Review Letters* (10.1103/PhysRevLett.105.176601 / Phys. Rev. Lett. 105, 176601 (2010)).

Press Office

Franziska Rott
Tel.: +49 (0)30-8062-42607
Fax: +49 (0)30-8062-42998
franziska.rott@helmholtz-berlin.de

Being made from carbon compounds, in other words plastics, organic solar cells are also known as plastic solar cells. The heart of the cell is a layer only a hundred millionth of a millimetre thick, made of two components, polymers and soccer ball-shaped fullerenes, mixed together. When light strikes a layer of this mixture, the polymer component is set into an excited state, dubbed an exciton. When an exciton bumps into a fullerene, an electron jumps over to the soccer ball molecule and a "hole" remains behind in the polymer. So that current can flow, the electrons and holes must travel to their respectively opposite contacts. The electrons travel via the fullerenes while the holes travel via the polymer chain. The holes, which scientists call polarons, can obstruct one another along their path and thus reduce the efficiency of the solar cell. This sets the limit on how much electrical energy can be yielded from a given amount of solar energy.

Using electrically detected magnetic resonance (EDMR), the scientists demonstrated that the polarons always get in one another's way when their magnetic moment (spin) is identical. "For the first time, we have uncovered and thus proven the long-assumed formation of these so-called bipolarons," says Jan Behrends, who performed the measurements during his doctorate at the HZB Institute for Silicon Photovoltaics.

The researchers' EDMR method involved manipulating the spin of the polarons using an external magnetic field and a microwave pulse. Using a resonance effect, the randomly distributed spin could be turned and aimed like a compass needle. Measurements revealed that current flows freely when the tiny magnets are oppositely aligned, but is blocked when they are aligned in the same direction.

The researchers demonstrated these current losses in plastic solar cells at room temperature, having redesigned an experimental method originally developed for silicon. “With this important finding, we should soon see advancements in organic solar cell technology as new plastics are introduced that develop no spin blockades”, says project leader Dr. Klaus Lips.

The **Helmholtz-Zentrum Berlin für Materialien und Energie (HZB)** operates and develops large scale facilities for research with photons (synchrotron beams) and neutrons. The experimental facilities, some of which are unique, are used annually by more than 2,500 guest researchers from universities and other research organisations worldwide. Above all, HZB is known for the unique sample environments that can be created (high magnetic fields, low temperatures). HZB conducts materials research on themes that especially benefit from and are suited to large scale facilities. Research topics include magnetic materials and functional materials. In the research focus area of solar energy, the development of thin film solar cells is a priority, whilst chemical fuels from sunlight are also a vital research theme. HZB has approx. 1,100 employees of whom some 800 work on the Lise-Meitner Campus in Wannsee and 300 on the Wilhelm-Conrad-Röntgen Campus in Adlershof. HZB is a member of the Helmholtz Association of German Research Centres, the largest scientific organisation in Germany.