



Cite this: *Dalton Trans.*, 2019, **48**, 11142

Correction: Magnetic relaxation in cobalt(II)-based single-ion magnets influenced by distortion of the pseudotetrahedral [N₂O₂] coordination environment

Michael Böhme,^a Sven Ziegenbalg,^a Azar Aliabadi,^b Alexander Schnegg,^b Helmar Görls^a and Winfried Plass^{*,a}

DOI: 10.1039/c9dt90143d

rsc.li/dalton

Correction for 'Magnetic relaxation in cobalt(II)-based single-ion magnets influenced by distortion of the pseudotetrahedral [N₂O₂] coordination environment' by Michael Böhme *et al.*, *Dalton Trans.*, 2018, **47**, 10861–10873.

In Fig. 4 and 5 the assignment of the complexes **1** and **2** to the graphs needs to be altered. The correct Fig. 4 and 5 for **1** (black) and **2** (red) are shown below (here denoted as Fig. 1 and 2, respectively).




Fig. 1 Relaxation times as the log–log plot of τ_c^{-1} vs. T for **1** and **2** with applied dc fields of 400 and 1000 Oe. Lines representing the slopes indicative of direct ($\sim T$) and Raman relaxation processes ($\sim T^5$ and $\sim T^9$) are depicted as a guide for the eye.⁹⁸

^aInstitut für Anorganische und Analytische Chemie, Friedrich-Schiller-Universität Jena, Humboldtstraße 8, 07745 Jena, Germany. E-mail: sekr.plass@uni-jena.de; Fax: +49 3641 948132; Tel: +49 3641 948130

^bBerlin Joint EPR Lab, Institute for Nanospectroscopy, Helmholtz-Zentrum Berlin für Materialien und Energie, Kekuléstraße 5, 12489 Berlin, Germany






Fig. 2 Arrhenius plot of relaxation times as $\ln \tau_c$ vs. $1/T$ for 1 and 2 with applied dc fields of 400 and 1000 Oe. Lines represent simulated values for best-fit parameters.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

